



Rewarding Learning

**ADVANCED**  
**General Certificate of Education**  
**2025**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

# Chemistry

Assessment Unit A2 3

*assessing*

Further Practical Chemistry

**Practical Booklet B (Theory)**



**[ACH32]**

\*ACH32\*

**FRIDAY 20 JUNE, MORNING**

## TIME

1 hour 15 minutes.

## INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

**You must answer the questions in the spaces provided.**

**Do not write outside the boxed area on each page or on blank pages.**

Complete questions in black ink and use a dark HB pencil for drawings and graphs.

**Do not write with a gel pen.**

Answer **all four** questions.

## INFORMATION FOR CANDIDATES

The total mark for this paper is 60.

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

You may use a scientific calculator.


A Data Leaflet, which includes a Periodic Table of the Elements, is included in this question paper.

14558



\*16ACH3201\*

- 1 Glucose,  $C_6H_{12}O_6$ , and other simple sugars are the monomers which form carbohydrates such as starch. The structure of glucose is shown below.



- (a) Explain why glucose is soluble in water but insoluble in hexane.

---

---

---

---

[2]

- (b) On heating in a boiling tube, glucose decomposes to form a black solid and droplets of a colourless liquid which are visible on the sides of the boiling tube.

- (i) Suggest a name for the colourless liquid.

---

[1]

- (ii) Suggest a name for the black solid.

---

[1]



- (c) Two experiments were carried out using an aqueous solution of glucose. The methods and observations for the two experiments are shown in the table below.

Experiment	Method	Observations
1	Add glucose solution to acidified potassium dichromate(VI) solution and heat the reaction mixture under reflux	orange solution changes to green
2	Add Tollens' reagent to glucose solution in a boiling tube and warm the contents of the boiling tube in a water bath	colourless solution changes to form a silver mirror

- (i) Draw a labelled diagram of the assembled apparatus required to heat the reaction mixture under reflux in Experiment 1.

[3]

[Turn over



- (ii) The half-equation below represents one of the possible oxidation reactions of glucose in Experiment 1.



Using the half-equation above, write a redox equation for the reaction occurring when glucose reacts with acidified dichromate(VI) ions.

\_\_\_\_\_ [2]

- (iii) Draw the structure of the oxidation product,  $\text{C}_6\text{H}_{10}\text{O}_8$ , formed in (c)(ii).

[2]

- (iv) Which functional group in glucose causes it to react with Tollens' reagent in Experiment 2?

\_\_\_\_\_ [1]



(v) What would be observed if the glucose solution was mixed with Fehling's solution and warmed in a water bath?

\_\_\_\_\_

\_\_\_\_\_ [2]

(d) Starch solution is used as an indicator in iodine-thiosulfate titrations.

(i) State the colour change observed at the end point when starch indicator is used in an iodine-thiosulfate titration.


From \_\_\_\_\_ to \_\_\_\_\_ [1]

(ii) Write an equation for the reaction of sodium thiosulfate with iodine.

\_\_\_\_\_ [2]



- 2 Phenylamine (density  $1.02 \text{ g cm}^{-3}$ ) reacts with ethanoyl chloride (density  $1.10 \text{ g cm}^{-3}$ ) to form the solid N-phenylethanamide. The equation for the reaction is given below.



- (a) Draw the skeletal formula of N-phenylethanamide.

[1]

- (b) A  $3.5 \text{ cm}^3$  sample of phenylamine was mixed with a  $1.5 \text{ cm}^3$  sample of ethanoyl chloride.

- (i) Calculate the number of moles of phenylamine in the  $3.5 \text{ cm}^3$  sample.

Answer \_\_\_\_\_ [2]

- (ii) Calculate the number of moles of ethanoyl chloride present in the  $1.5 \text{ cm}^3$  sample.

Answer \_\_\_\_\_ [2]





(ii) Describe, giving practical details, how the melting point of the solid N-phenylethanamide could be determined.

---

---

---

---

---

[3]

(iii) State two ways in which the melting point of an impure sample of N-phenylethanamide would differ from the melting point of a pure sample.

1. \_\_\_\_\_

---

2. \_\_\_\_\_

---

[2]





**BLANK PAGE**

**DO NOT WRITE ON THIS PAGE**

**(Questions continue overleaf)**

14558

**[Turn over**



\*16ACH3209\*

3 Solid **X** is a hydrated iron(III) compound.

(a) Solid **X** dissolves in water to form a yellow solution. Write the formula of the hexaaqua complex ion present in the yellow solution.

\_\_\_\_\_ [1]

(b) On addition of sodium hydroxide solution to an aqueous solution of **X**, a brown precipitate is formed. Write an equation for the reaction that occurs representing the brown precipitate as  $[\text{Fe}(\text{OH})_3(\text{H}_2\text{O})_3]$ .

\_\_\_\_\_ [1]

(c) When acidified silver nitrate solution is added to an aqueous solution of **X**, a white precipitate is formed. The white precipitate dissolves on addition of concentrated ammonia solution.

(i) Name the anion present in the aqueous solution of **X**.

\_\_\_\_\_ [1]

(ii) The silver ions in the white precipitate react with ammonia solution to form a soluble complex which has a linear shape. Suggest the formula of this complex.

\_\_\_\_\_ [1]

(d) When a spatula-measure of sodium carbonate is added to an aqueous solution of **X**, bubbles of gas are observed.

Identify the gas produced and suggest what the formation of this gas indicates about the aqueous solution of **X**.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_ [2]



- (e) The addition of a solution containing thiocyanate ions ( $\text{SCN}^-$ ) to an aqueous solution of **X** resulted in the formation of a blood red solution. This is a ligand replacement reaction in which one thiocyanate ion replaces one molecule of water in the hexaqua iron(III) ion. Suggest an equation for the reaction that occurs.

\_\_\_\_\_ [1]

- (f) 1 mole of the hydrated iron(III) compound contains 6 moles of water of crystallisation. The percentage of water of crystallisation in the compound is 39.93%.

- (i) Calculate the relative formula mass of the hydrated iron(III) compound. Give your answer to 1 decimal place.

Answer \_\_\_\_\_ [2]

- (ii) Based on your calculation in (f)(i) and the information in (a) to (e), write the formula of the hydrated iron(III) compound.

\_\_\_\_\_ [1]

[Turn over



4 A method for a back titration is outlined below.

**Step 1:** Weigh out approximately 2.5 g of a Group II metal oxide in a beaker.

**Step 2:** Add 50.0 cm<sup>3</sup> of 2.00 mol dm<sup>-3</sup> hydrochloric acid to the Group II metal oxide in the beaker.

**Step 3:** Make up the reaction mixture from **Step 2** to form a 250 cm<sup>3</sup> aqueous solution in a volumetric flask.

**Step 4:** Titrate 25.0 cm<sup>3</sup> samples of the solution from **Step 3** against 0.100 mol dm<sup>-3</sup> sodium hydroxide solution using a suitable indicator.

(a) (i) Define the term **back titration**.

---

---

---

[2]

(ii) Describe the practical procedure carried out in **Step 3** of the method.

---

---

---

---

---

---

---

---

[4]

(iii) Name a suitable indicator for the titration in **Step 4**. State the colour change at the end point of this titration.

---

---

[2]



(b) 2.44 g of a Group II metal oxide were used in the back titration and the results obtained are shown in the table below.

Titration	Initial burette reading /cm <sup>3</sup>	Final burette reading /cm <sup>3</sup>	Titre /cm <sup>3</sup>
1	0.0	13.3	13.3
2	13.3	26.2	12.9
3	26.2	39.0	12.8

(i) Explain why the titre value from **Titration 1** should not be used in the calculation of the mean titre.

\_\_\_\_\_ [1]

(ii) Calculate the number of moles of hydrochloric acid added to the Group II metal oxide in **Step 2**.

Answer \_\_\_\_\_ [1]

(iii) Calculate the number of moles of hydrochloric acid neutralised by the 0.100 mol dm<sup>-3</sup> sodium hydroxide solution in **Step 4**.

Answer \_\_\_\_\_ [2]

(iv) Using MO to represent the Group II metal oxide, write an equation for the reaction of the metal oxide with hydrochloric acid.

\_\_\_\_\_ [1]

[Turn over



- (v) Calculate the relative formula mass of the Group II metal oxide, MO.  
Give your answer to the nearest whole number.

Answer \_\_\_\_\_ [3]

---

**THIS IS THE END OF THE QUESTION PAPER**

---





**BLANK PAGE**  
**DO NOT WRITE ON THIS PAGE**

14558



\*16ACH3215\*

**DO NOT WRITE ON THIS PAGE**

For Examiner's use only		
Question Number	Examiner Mark	Remark
1		
2		
3		
4		
<b>Total Marks</b>		

Permission to reproduce all copyright material has been applied for.  
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA will be happy to rectify any omissions of acknowledgement in future if notified.

ACH32/5  
297934



\*16ACH3216\*



## General Information

1 tonne =  $10^6$  g

1 metre =  $10^9$  nm

One mole of any gas at 293 K and a pressure of 1 atmosphere ( $10^5$  Pa) occupies a volume of 24 dm<sup>3</sup>

Avogadro Constant =  $6.02 \times 10^{23}$  mol<sup>-1</sup>

Planck Constant =  $6.63 \times 10^{-34}$  Js

Specific Heat Capacity of water =  $4.2 \text{ J g}^{-1} \text{ K}^{-1}$

Speed of Light =  $3 \times 10^8 \text{ ms}^{-1}$



## Characteristic absorptions in IR spectroscopy

Wavenumber/cm <sup>-1</sup>	Bond	Compound
550–850	C–X (X = Cl, Br, I)	Haloalkanes
750–1100	C–C	Alkanes, alkyl groups
1000–1300	C–O	Alcohols, esters, carboxylic acids
1450–1650	C=C	Arenes
1600–1700	C=C	Alkenes
1650–1800	C=O	Carboxylic acids, esters, aldehydes, ketones, amides, acyl chlorides
2200–2300	C≡N	Nitriles
2500–3200	O–H	Carboxylic acids
2750–2850	C–H	Aldehydes
2850–3000	C–H	Alkanes, alkyl groups, alkenes, arenes
3200–3600	O–H	Alcohols
3300–3500	N–H	Amines, amides

## Proton Chemical Shifts in Nuclear Magnetic Resonance Spectroscopy (relative to TMS)

Chemical Shift	Structure	
0.5–2.0	–CH	Saturated alkanes
0.5–5.5	–OH	Alcohols
1.0–3.0	–NH	Amines
2.0–3.0	–CO–CH	Ketones
	–N–CH	Amines
	C <sub>6</sub> H <sub>5</sub> –CH	Arene (aliphatic on ring)
2.0–4.0	X–CH	X = Cl or Br (3.0–4.0) X = I (2.0–3.0)
4.5–6.0	–C=CH	Alkenes
5.5–8.5	RCONH	Amides
6.0–8.0	–C <sub>6</sub> H <sub>5</sub>	Arenes (on ring)
9.0–10.0	–CHO	Aldehydes
10.0–12.0	–COOH	Carboxylic acids

These chemical shifts are concentration and temperature dependent and may be outside the ranges indicated above.

© CCEA 2017

COUNCIL FOR THE CURRICULUM, EXAMINATIONS AND ASSESSMENT

29 Clarendon Road, Clarendon Dock, Belfast BT1 3BG

Tel: +44 (0)28 9026 1200 Fax: +44 (0)28 9026 1234

Email: info@ccea.org.uk Web: www.ccea.org.uk



# Data Leaflet

## Including the Periodic Table of the Elements

For the use of candidates taking  
Advanced Subsidiary and  
Advanced Level Examinations

**Copies must be free from notes or additions of any kind. No other type of data booklet or information sheet is authorised for use in the examinations**

# gce a/as examinations chemistry

