



General Certificate of Secondary Education
November 2022

Centre Number

| | | | | |
|--|--|--|--|--|
| | | | | |
|--|--|--|--|--|

Candidate Number

| | | | | |
|--|--|--|--|--|
| | | | | |
|--|--|--|--|--|

Mathematics

Unit M7 Paper 1
(Non-Calculator)

Higher Tier



[GMC71]

GMC71

THURSDAY 1 DECEMBER, 9.15am–10.30am

TIME

1 hour 15 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page. **You are provided with Higher Tier Additional Support Materials for use with this paper.**

You must answer the questions in the spaces provided.

Do not write outside the boxed area on each page, on blank pages or tracing paper.

Complete in black ink only. **Do not write with a gel pen.**

Answer **all seventeen** questions.

All working should be clearly shown in the spaces provided. Marks may be awarded for partially correct solutions.

You **must not** use a calculator for this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 50.

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

You should have a ruler, compasses and a protractor.

The Formula Sheet is on page 2.

13333



20GMC7101

Formula Sheet

Volume of prism = area of cross section \times length



Area of trapezium = $\frac{1}{2}(a+b)h$



Volume of sphere = $\frac{4}{3}\pi r^3$

Surface area of sphere = $4\pi r^2$




Volume of cone = $\frac{1}{3}\pi r^2 h$

Curved surface area of cone = $\pi r l$



In any triangle ABC



Quadratic Equation

The solutions of $ax^2 + bx + c = 0$
where $a \neq 0$, are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$


Sine Rule: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

Cosine Rule: $a^2 = b^2 + c^2 - 2bc \cos A$

Area of triangle = $\frac{1}{2} ab \sin C$



1 A sequence consists of rectangles and circles as shown.



(a) Complete the table for patterns 2, 3, 4 and 7

| | Pattern 1 | Pattern 2 | Pattern 3 | Pattern 4 | Pattern 7 |
|--|-----------|-----------|-----------|-----------|-----------|
| Number of rectangles | 1 | 2 | | | |
| Number of circles | 4 | | 8 | | |
| Total number of rectangles and circles | 5 | | | | |

[2]


(b) Which pattern number has a total number of 38 rectangles and circles?

Answer Pattern number _____ [1]

[Turn over



2 The distance–time graph shows a train journey.



(a) What is the distance between Bangor and Lisburn?

Answer _____ miles [1]

(b) Give a reason for the horizontal line which starts when the train reaches Lisburn.

Answer _____
_____ [1]

(c) The train leaves Lisburn and travels on to Lurgan.

The distance between Lisburn and Lurgan is 13 miles and the journey takes 15 minutes.

Show this information on the graph. [1]



(d) What is the average speed of the train from Lisburn to Lurgan?

Answer _____ mph [1]

(e) Give one reason for knowing that the graph shows average speed rather than actual speed.

Answer _____ [1]

3 Ross works out $200 \times 90 = 18\,000$

Write down a calculation Ross could use to check his answer.

Answer _____ [1]

[Turn over



4 Calculate 5% of 45


Answer _____ [2]

13333



20GMC7106

5 Charlie has drawn a **sketch** of a trapezium with lengths and angles as shown.



AB is already drawn.

Using a ruler and protractor, complete Charlie's trapezium **accurately**.



[4]

[Turn over



- 6 (a) 60% of entrants pass a test.

What is the ratio of

number who pass : number who fail?

Give your answer in simplest form.

Answer _____ [2]

- (b) The ratio of boys : girls born in a hospital one week was 3 : 5

What fraction were girls?

Answer _____ [1]

- 7 At the beginning of January an oil tank was $\frac{3}{4}$ full.

At the end of March it was $\frac{1}{6}$ full.


What fraction of the tank was used?

Show your working.

Answer _____ [2]



8



(a) Draw the image of shape A after a translation, 7 right and 4 down.

Label it T.

[1]

(b) Draw the image of shape A after a rotation, 90° anticlockwise, about the origin.


Label it R.

[2]


[Turn over




9



Quadrilateral



Pentagon



Hexagon

P is a polygon.

The sum of all the angles in P is three times the sum of the angles in a quadrilateral.

How many sides has P?

Answer _____ [2]



10 A shape has a perimeter of 16 cm and an area of 10 cm^2

It is enlarged using a scale factor of 2

(a) What is the perimeter of the enlarged shape?

Answer _____ cm [1]

(b) How many times bigger is the area of the enlarged shape than the area of the original shape?

Answer _____ [1]



11 Each interior angle of a regular polygon is 140° .

How many sides has the polygon?

Answer _____ [2]




12 The diagram shows a section of coastline with a lifeboat station marked at B and a lighthouse marked at L.

A sinking ship sends a distress signal.

The ship is less than 70 km from B and less than 30 km from L.

Using a scale of 1 cm = 10 km, **shade the region** in which the ship could be.



[3]

[Turn over

13333



20GMC7113


13 (a) Complete the table for $y = x^2 + 3x - 3$

| | | | | | | | |
|-----|----|----|----|----|----|---|---|
| x | -4 | -3 | -2 | -1 | 0 | 1 | 2 |
| y | 1 | | -5 | -5 | -3 | 1 | |

[2]

(b) Draw the graph of $y = x^2 + 3x - 3$ from $x = -4$ to $x = 2$

[2]



(c) Use your graph to estimate the solutions to $x^2 + 3x - 3 = 0$

Answer _____ [2]



14 (a) Write the decimal number 15 as a binary number.

Answer _____ [1]

(b) Write the binary number 1000000 as a decimal number.

Answer _____ [1]



15 An industrial tank has dimensions 10 times larger than the dimensions of a similar domestic tank.

How many times larger is its volume?

Answer _____ [1]



16 Tom has some 20p coins and some 50p coins.

The total value of Tom's coins is £8.60

Seana has twice as many 20p coins as Tom, but only half as many 50p coins as Tom.

The total value of Seana's coins is £8.20

Find how many of each type of coin Tom has.

A solution by trial and improvement will not be accepted.

Answer _____ 20p coins, _____ 50p coins [5]


[Turn over

13333



20GMC7117

17 (a)




The lines $y = 2x + 1$ and $x + y = 4$ have been drawn on the grid.

By drawing another line on the grid above, indicate clearly by the letter R

the region satisfying $y \geq 2x + 1$ and $x + y \leq 4$ and $x \geq -1$ [2]



(b)



By drawing more lines on the grid above, indicate clearly by the letter B the region satisfying $y \leq 2x$ and $x + y \leq 4$ and $y \geq 1$ [2]

THIS IS THE END OF THE QUESTION PAPER



Sources: All images © CCEA unless stated

DO NOT WRITE ON THIS PAGE

| For Examiner's use only | |
|-------------------------|-------|
| Question Number | Marks |
| 1 | |
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| 6 | |
| 7 | |
| 8 | |
| 9 | |
| 10 | |
| 11 | |
| 12 | |
| 13 | |
| 14 | |
| 15 | |
| 16 | |
| 17 | |

| | |
|--------------------|--|
| Total Marks | |
|--------------------|--|

Examiner Number

Permission to reproduce all copyright material has been applied for.
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA will be happy to rectify any omissions of acknowledgement in future if notified.

13333/5



20GMC7120



Rewarding Learning

**General Certificate of Secondary Education
November 2022**

GCSE Mathematics

HIGHER TIER ADDITIONAL SUPPORT MATERIALS (For use in November 2022)

HIGHER TIER ADDITIONAL SUPPORT MATERIALS (November 2022)

Numbers

Highest Common Factor (HCF): The highest common factor is the largest number that divides evenly into two or more numbers, e.g. 4 is the HCF of 8 and 20

Trial and Improvement

This is a method of trying different values in an equation until you get a suitable solution (e.g. to 1 decimal place).


Range

The range of a set of data is the difference between the largest value and the smallest value in the data set.

Mean

The mean of a set of data is the sum of all the data values divided by the number of data values.

Area and Volume



The area of a rectangle is found by multiplying the length of the rectangle by the breadth.

$A = L \times B$ where A is area, L is length and B is breadth.

The area of a triangle is found by multiplying half the length of the base by the perpendicular height of the triangle.

$A = \frac{1}{2}bh$ where b is the base and h is the perpendicular height of the triangle.

The area of a circle is $A = \pi r^2$ where r is the radius of the circle.

The volume of a cuboid is found by multiplying the length by the breadth by the height of the cuboid.

$V = L \times B \times H$ where V is volume, L is length, B is breadth and H is height.

Angles

There are 180° on a straight line.

There are 180° inside a triangle.

An isosceles triangle is a triangle with 2 equal sides and 2 equal angles.

The sum of all the angles inside a polygon is given by $180(n - 2)$ where n is the number of sides in the polygon.

Pie Chart

In a pie chart, the total angle that corresponds to the entire data set is 360°

Probability

The sum of the probabilities of all outcomes equals 1

Compound Measures


$$\text{Average Speed} = \frac{\text{Distance}}{\text{Time}}$$

$$\text{Density} = \frac{\text{Mass}}{\text{Volume}}$$

Pythagoras' Theorem

If a , b and c are the sides of a right angled triangle shown below, then

$$a^2 + b^2 = c^2$$



Midpoint of a line

If (x_1, y_1) and (x_2, y_2) are the end points of a line, then the coordinates of the midpoint M of the line are

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Gradient of a line

If (x_1, y_1) and (x_2, y_2) are two points on a line, then the gradient m of the line is

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

Lines

Parallel lines have the same gradient.

If a straight line has gradient m , then a line which is perpendicular to this line has a gradient $-\frac{1}{m}$

Equation of a circle

The equation of a circle of radius r and centre, the origin $(0, 0)$, is given by $x^2 + y^2 = r^2$ where (x, y) are the coordinates of any point on the circumference of the circle.

Tangent/Radius property

The tangent to a circle is perpendicular to the radius at the point of contact with the circle.


Cyclic Quadrilateral property

The opposite angles of a cyclic quadrilateral add up to 180°

Alternate Segment Theorem

In a circle, the angle between a chord and a tangent through one of the end points of the chord is equal to the angle in the alternate segment.


Trigonometric ratios in right angled triangles



$$\sin x^\circ = \frac{O}{H} \quad \cos x^\circ = \frac{A}{H} \quad \tan x^\circ = \frac{O}{A}$$

Angle of depression

If a person stands and looks down at an object, the **angle of depression** is the angle between the horizontal line of sight and the object.



Frequency density in histograms

$$\text{Frequency density} = \frac{\text{Frequency}}{\text{Class width}}$$