



Rewarding Learning

General Certificate of Secondary Education
2023

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Mathematics

Unit M7 Paper 1
(Non-Calculator)

Higher Tier



[GMC71]

GMC71

WEDNESDAY 7 JUNE, 9.15 am–10.30 am

TIME

1 hour 15 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page. **You are provided with Higher Tier Additional Support Materials for use with this paper.**

You must answer the questions in the spaces provided.

Do not write outside the boxed area on each page, on blank pages or tracing paper.

Complete in black ink only. **Do not write with a gel pen.**

Answer **all thirteen** questions.

All working should be clearly shown in the spaces provided. Marks may be awarded for partially correct solutions.

You **must not** use a calculator for this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 50.

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

You should have a ruler, compasses and a protractor.

The Formula Sheet is on page 2.

13346.08 R



16GMC7101

Formula Sheet

Volume of prism = area of cross section \times length



Area of trapezium = $\frac{1}{2}(a+b)h$



Volume of sphere = $\frac{4}{3}\pi r^3$

Surface area of sphere = $4\pi r^2$




Volume of cone = $\frac{1}{3}\pi r^2 h$

Curved surface area of cone = $\pi r l$



In any triangle ABC



Quadratic Equation

The solutions of $ax^2 + bx + c = 0$
where $a \neq 0$, are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Sine Rule: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

Cosine Rule: $a^2 = b^2 + c^2 - 2bc \cos A$

Area of triangle = $\frac{1}{2} ab \sin C$



1 There are 30 pupils in a class.

$\frac{3}{5}$ are girls.

$\frac{2}{9}$ of the girls are absent.

25% of the boys are absent.

(a) Complete the table to show this information.

	Girls	Boys
Number of pupils		
Absent		
Present		

[2]

(b) What fraction of the pupils are present?

Answer _____ [1]

[Turn over

13346.08 R



16GMC7103

2 The pulse rate of an athlete is taken before and after training.

His starting pulse was 54 beats per minute.

After training, it increased by $\frac{1}{6}$ of this value.


Show that each beat now lasts less than 1 second.




[4]



3 Two fair spinners are each spun once.



Spinner A



Spinner B

The scores on each spinner are multiplied together.

(a) Complete the table of outcomes.

		Spinner A			
		10	20	30	40
Spinner B	1				
	2				
	3				

[2]

(b) What is the probability that the outcome is 60?

Answer _____ [1]

(c) What is the probability that the outcome is less than 40?

Answer _____ [2]

[Turn over



4 The probability that a child is left handed is q .

What is the probability that a child is not left handed?

Answer _____ [1]

5 (a) Given that $23 \times 146 = 3358$

write down the answer to 2.3×1.46

Answer _____ [1]

(b) Showing clearly how you do it, **estimate** the answer to

$$\frac{202 \times 29}{0.48}$$

Answer _____ [3]



6 In April, 30 males and 20 females took their driving test.
50% of the males passed and 40% of the females passed.
What percentage of people who took the test failed?




Answer _____ % [3]

[Turn over

13346.08 R



16GMC7107




- (a) Describe fully the single transformation that maps shape A onto shape B.

Answer _____ [2]




(b) A sketch of a parallelogram PQRS is shown.



It is not drawn to scale.

Use a ruler and protractor to draw an accurate diagram of the parallelogram in the box below.



[4]

[Turn over



8 Tony owns a pizza takeaway.

Every time he sells 100 pizzas, he records the number that were pepperoni.

The table shows some of his data.

Number of pizzas sold	Number of pepperoni pizzas	Relative frequency of pepperoni pizzas
100	17	0.17
100	23	0.2
100	23	0.21
100	25	
100		0.23

(a) Complete the table.

[4]

(b) Tony uses his data to predict next year's pizza sales.

He thinks he will sell 2300 pepperoni pizzas next year.

How many pizzas is he expecting to sell in total?


Answer _____ [1]



9 By drawing a suitable line on the grid, solve the simultaneous equations

$$x - 2y = 4$$

$$y = 3x + 3$$



Answer $x =$ _____ $y =$ _____ [4]

[Turn over

13346.08 R



16GMC7111

10 (a) The first four terms of a sequence are

1, 4, 7, 10

What is the n^{th} term for this sequence?

Answer _____ [2]

(b) Hence find the n^{th} term for the sequence below.

$\frac{1}{1}$, $\frac{4}{4}$, $\frac{9}{7}$, $\frac{16}{10}$

Answer _____ [2]

11 In standard form

$$(2 \times 10^x) \times (3 \times 10^y) = 6 \times 10^{11}$$

$$(3 \times 10^y) \div (2 \times 10^x) = 1.5 \times 10^3$$



Find the values of x and y .

Show your working.

Answer $x =$ _____ $y =$ _____ [4]



12



The semicircle and the triangle have the same perimeter.

Find the exact value of r , giving your answer in terms of π .

Answer $r =$ _____ [4]



13 (a) A dice is rolled once and a coin is flipped once.

Here are 4 events.

Event A – the roll of the dice is even

Event B – the roll of the dice is odd

Event C – the flipped coin lands heads

Event D – the flipped coin lands tails

(i) Write down two of these events that are independent.

Answer Event _____ and Event _____ [1]

(ii) Write down two of these events that are mutually exclusive.

Answer Event _____ and Event _____ [1]

(b) Two events, P and Q, are mutually exclusive.

What is the probability that either event P or event Q occurs?

Circle your answer.

Probability(P) + Probability(Q)

Probability(P) × Probability(Q)

more information is needed

[1]





THIS IS THE END OF THE QUESTION PAPER

BLANK PAGE

DO NOT WRITE ON THIS PAGE

13346.08 R



16GMC7115

Sources: All images © CCEA unless stated

DO NOT WRITE ON THIS PAGE

For Examiner's use only	
Question Number	Marks
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	

Total Marks	
--------------------	--

Examiner Number

Permission to reproduce all copyright material has been applied for.
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA will be happy to rectify any omissions of acknowledgement in future if notified.

13346.08 R



16GMC7116



Rewarding Learning

**General Certificate of Secondary Education
Summer 2023**

GCSE Mathematics


HIGHER TIER ADDITIONAL SUPPORT MATERIALS (For use in Summer 2023)

HIGHER TIER ADDITIONAL SUPPORT MATERIALS (Summer 2023)

$$\text{Average Speed} = \frac{\text{Distance}}{\text{Time}}$$

Perimeter, Area and Volume

The perimeter of a polygon is the distance around the outside of the polygon.



The area of a rectangle is found by multiplying the length of the rectangle by the breadth.

$A = L \times B$ where L is length and B is breadth.

The volume of a cuboid is found by multiplying the length by the breadth by the height of the cuboid.

$V = L \times B \times H$ where V is volume, L is length, B is breadth and H is height.

The area of a circle is $A = \pi r^2$ where r is the radius of the circle.

The circumference (perimeter) of a circle is $C = 2\pi r$ where r is the radius of the circle. An alternative formula is $C = \pi d$ where d is the diameter of the circle.

Mid point of a line

If (x_1, y_1) and (x_2, y_2) are the end points of a line, then the coordinates of the midpoint M of the line are

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Gradient of a line

If (x_1, y_1) and (x_2, y_2) are two points on a line, then the gradient m of the line is

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

Lines

Parallel lines have the same gradient.

If a straight line has gradient m , then a line which is perpendicular to this line has a gradient $-\frac{1}{m}$

Geometry and Angles

There are 180° on a straight line.


There are 180° inside a triangle.

An isosceles triangle is a triangle with 2 equal sides and 2 equal angles.

The sum of all the angles inside a polygon is given by $180(n - 2)$ where n is the number of sides in the polygon.

Angle of elevation


If a person stands and looks up at an object, the **angle of elevation** is the angle between the horizontal line of sight and the object.




Pythagoras' Theorem

If a , b and c are the sides of a right angled triangle shown below, then

$$a^2 + b^2 = c^2$$



Trigonometric ratios in right angled triangles



$$\sin x^\circ = \frac{O}{H} \quad \cos x^\circ = \frac{A}{H} \quad \tan x^\circ = \frac{O}{A}$$

Tangent/Radius property

The tangent to a circle is perpendicular to the radius at the point of contact with the circle.

Alternate Segment Theorem

In a circle, the angle between a chord and a tangent through one of the end points of the chord is equal to the angle in the alternate segment.

Mean

The mean of a set of data is the sum of all the data values divided by the number of data values.

Estimate for the mean of a grouped frequency distribution

Estimated mean = sum of (mid interval values multiplied by their frequency) divided by the sum of all the frequencies.

Pie Chart

In a pie chart, the total angle that corresponds to the entire data set is 360°

Probability

The sum of the probabilities of all outcomes equals 1

Frequency density in histograms

$$\text{Frequency density} = \frac{\text{Frequency}}{\text{Class width}}$$

